



DAVID B. COHEN, M.D.

84 North Main Street Bld#2 Branford, CT 06405 203.483.2024

CALL OUR OFFICE AT 203.407.3518 FOR FOLLOW-UP OR QUESTIONS

DISCHARGE/HOME INSTRUCTIONS
Achilles Tendon Repair

Patient Name:

Follow up appointment:

- Call your doctor later today or tomorrow to arrange to be seen in 10 - 14 days if you do not already have one.
- COS Surgical Center wishes you a **speedy recovery**. Please note the following guidelines for your recovery period at home:

DIET:

- Your regular diet may be resumed today. Start with liquids. Drink additional fluids today.
- Avoid alcohol for 24 hours and while you are taking your pain pills.

MEDICATIONS: It is normal to experience discomfort after surgery. Take medication as prescribed.

- Pain medication may cause constipation. Use an over-the-counter stool softener (ie. Colace) as needed.
- Eat something before taking medication to avoid stomach upset.
- Your pain medication is Oxycodone. Take 1-2 when you are aware of soreness and then every 3-4 hours as needed. Set an alarm clock to wake you up at night to take the medicine during the first few days. It takes 45 minutes to start working.
- Use Naprosyn (an anti-inflammatory) for 14 days after surgery. Take one pill in the morning, one in the evening with food. Discontinue use if you have any history of ulcer disease or difficulty using anti-inflammatory medications or aspirin.

DRESSING: KEEP YOUR DRESSING DRY AND CLEAN.

- Keep ankle splint dry.

ACTIVITY:

- Do not drive, operate machinery, or sign legal documents. You may be forgetful due to today's medications.
- Rest today. Get up every 2-3 hours during the day. You may experience light-headedness and sleepiness following your surgery. We encourage you to have someone with you for 24 hours.
- To bathroom with assistance. Apply ice for minimum one week while awake (30 minutes on, 30 minutes off). You should use it after physical therapy as well.



- You may shower on post-operative day 3 but keep the splint dry. Cover with a plastic bag or cast guard (from pharmacy)
- Flex and extend other ankle to promote blood flow. Your ankle may swell when you are up and about. This is normal.
- Elevate leg for 4-5 days. Keep your leg elevated with a pillow under your calf, NOT under the knee. Keep your “toes above your nose.”
- Use crutches with NO weight bearing for 6 weeks.

NOTIFY YOUR SURGEON IF YOU ARE UNSURE OF YOUR PROGRESS OR IF:

- Dressing is too tight.
- Fever above 101°F
- Excessive swelling
- Continued nausea/vomiting
- Uncontrolled pain
- Unable to urinate
- Excessive bleeding
- Infection (red, drainage and/or sore)
- If coldness, numbness or change in color of fingers/toes or skin (hand may be numb for 24 hours after nerve block)

IF YOU CANNOT REACH YOUR DOCTOR AND NEED HELP OR ADVICE RIGHT AWAY, GO TO THE EMERGENCY ROOM NEAREST YOU. IN THE EVENT OF A TRUE EMERGENCY CALL 911.

I (Patient/Escort) have read these instructions. I understand them and have no questions at this time.

_____, M.D. _____
Patient / Escort

MD OFFICE NUMBERS

Branford (203) 483.2509	Milford (203) 882.3373
Guilford (203) 453.6340	Orange (203) 795.4784
Hamden (203) 407.3500	Wallingford (203) 265.1800